

CHOW GROUPS AND HIGHER CONGRUENCES FOR THE NUMBER OF RATIONAL POINTS ON PROPER VARIETIES OVER FINITE FIELDS

NAJMUDDIN FAKHRUDDIN

ABSTRACT. Given a proper family of varieties over a smooth base, with smooth total space and general fibre, all over a finite field k with q elements, we show that a finiteness hypotheses on the Chow groups, CH_i , $i = 0, 1, \dots, r$, of the fibres in the family leads to congruences mod q^{r+1} for the number of rational points in *all* the fibres over k -rational points of the base. These hypotheses on the Chow groups are expected to hold for families of low degree intersections in many Fano varieties leading to a broad generalisation of the theorem of Ax–Katz [9], as well as results of the author and C. S. Rajan [7]. As an unconditional application, we give an asymptotic generalisation of the Ax–Katz theorem to low degree intersections in a large class of homogenous spaces.

1. INTRODUCTION

Let X be a smooth proper geometrically connected variety over a finite field k with q elements and assume that the base change maps of *rational* Chow groups, $CH_i(X) \rightarrow CH_i(X_{\overline{k}(X)})$ are isomorphisms for $i = 0, 1, \dots, r$ and some $r \geq 0$. These isomorphisms give rise to a decomposition of the diagonal correspondence in $X \times X$ (see Lemma 2.1 for details), which along with the Grothendieck–Lefschetz trace formula shows that $|X(k)| \equiv \sum_{i=0}^r b_{2i} q^i \pmod{q^{r+1}}$, where $b_i = \dim H_{\text{et}}^i(X_{\overline{k}}, \mathbb{Q}_\ell)$ with ℓ a prime not equal to $\text{char}(k)$; this was first observed by H. Esnault [4]. The goal of this paper is to prove similar results in the case where X is not necessarily smooth: that some such results should hold under fairly general hypotheses is suggested by the Ax–Katz theorem [9] which gives congruences as above for intersections of low degree hypersurfaces in $\mathbb{P}^n$. For $r = 0$, such results were proved by the author and C. S. Rajan in [7] under very general hypotheses; here we will show that a refinement of those methods implies congruences for singular varieties also when $r > 0$. Our results can thus be viewed as a motivic explanation for, as well as a broad generalisation of (albeit with some extra hypotheses which should conjecturally always hold), the theorem of Ax and Katz.

For singular varieties, even for $r = 0$, the Chow groups do not “control” congruences. The main observation of [7] was that if we consider a family of varieties over a smooth base with smooth total space, then CH_0 of the generic fibre (over a sufficiently large extension of the base field) does in fact “control” congruences mod q for the number of rational points of all closed fibres. If $r > 0$ it is easily seen, for example by blowing up a subvariety of the total space not intersecting the generic fibre, that a condition solely in terms of the Chow groups of the generic fibre cannot suffice to imply congruences mod q^{r+1} for closed fibres; some extra geometric and/or cohomological conditions must be added. The main result of this paper is then the following:

Theorem 1.1. *Let $f : X \rightarrow Y$ be a proper dominant morphism of smooth irreducible varieties with smooth geometrically irreducible generic fibre Z over a finite field k with q*

elements. For each point y of Y , let $Z_y = f^{-1}(y)$ and let $\mathcal{Y} = \text{Spec}(\mathcal{O}_{y,Y})$. Assume that for each $i = 0, 1, \dots, r$, the specialisation maps ([8, p.399])

$$CH_i(X \times_Y (\mathcal{Y} - \{y\})/\mathcal{Y} - \{y\}) \rightarrow CH_i(Z_y)$$

are surjective and the maps $CH_i(Z_y) \rightarrow CH_i((Z_y)_K)$ are isomorphisms for some embedding $k(y) \hookrightarrow K$ with K a universal domain and $i = 0, 1, \dots, r$. Further, assume that $R^j f_* \mathbb{Q}_\ell$ is a constant local system on Y for $j = 0, 1, \dots, 2r$. Then for all $y \in Y(k)$,

$$|Z_y(k)| \equiv \sum_{i=0}^r b_{2i} q^i \pmod{q^{r+1}},$$

where $b_{2i} = \text{rank } R^{2i} f_* \mathbb{Q}_\ell$.

Some condition similar to ours on the $R^j f_* \mathbb{Q}_\ell$ probably follows from the conclusion of the theorem (for all finite extensions of k); it appears to be close to the minimal extra condition not pertaining to the generic fibre needed to obtain congruences mod q^{r+1} for $r > 0$. But it seems likely that the hypotheses on the Chow groups of non-generic fibres is implied by the above cohomological condition along with the hypothesis on the Chow groups of the generic fibre.

A theorem of K. Paranjape [11] (which was improved in [6]) implies that the hypotheses of the theorem on Chow groups (for a fixed r) hold for the universal family of intersections of hypersurfaces of multidegree $(d_1, d_2, \dots, d_t)$ in $\mathbb{P}^n$ if n is sufficiently large compared to $\sum_{k=1}^t d_k$; the other conditions are easily checked. Thus we recover an asymptotic version of the Ax–Katz theorem and the full Ax–Katz theorem would follow from standard conjectures relating the coniveau filtration on cohomology and Chow groups. By partly extending Paranjape’s results to homogenous spaces we are able to obtain the following unconditional result:

Theorem 1.2. *Let k be finite field with q elements, let G_n be a k -form of one of the simple algebraic groups SL_{n+1} , Sp_{2n} , SO_{2n} or SO_{2n+1} and let P_n be a maximal parabolic subgroup of G_n also defined over k except for the case where G_n is Sp_{2n} and P_n is the stabilizer of a maximal isotropic subspace. Let $X_n = G_n/P_n$ and L_n the ample generator of $\text{Pic}(X_n)$. For any tuple $D := (d_1, d_2, \dots, d_t)$ of positive integers and any positive integer r , there exists an integer $N(D, r)$ such that for any $n > N(D, r)$ and any $s_i \in H^0(X_n, L_n^{\otimes d_i})$, $i = 1, 2, \dots, t$*

$$|Z(s_1, s_2, \dots, s_t)(k)| \equiv \sum_{j=0}^{r-1} \dim(b_2(X_n)) q^j \pmod{q^r}$$

where $Z(s_1, s_2, \dots, s_t)$ is the subscheme of X_n given by the vanishing of all the s_i ’s.

We also include an example (4.4) of a homogenous space for the exceptional group E_8 for which one also has “higher congruences” for the number of rational points on certain hypersurfaces.

The main idea of the proof of Theorem 1.1 is to use the hypothesis on the Chow groups to obtain a suitable decomposition of the diagonal correspondence in $X \times_Y X$ which, using standard properties of étale cohomology, implies the divisibility of the eigenvalues of Frobenius acting on a certain quotient of $H_c^*(X, \mathbb{Q}_\ell)$. An application of the Grothendieck–Lefschetz trace formula completes the proof. The decomposition is obtained by applying Lemma 2.1 to the generic fibre which gives us the desired decomposition except for an “error term” which

is a correspondence whose support does not intersect the generic fibre. This correspondence is controlled by applying Lemma 2.1 inductively to certain non-generic fibres along with a “nilpotence” argument which eventually shows that the error term can be neglected.

Notations and Conventions. By a variety we shall mean a reduced separated scheme of finite type over a field k . For X a variety, we shall abuse notation and denote by $H^*(X, \mathbb{Q}_\ell)$ the étale cohomology groups $H_{\text{ét}}^*(X_{\bar{k}}, \mathbb{Q}_\ell)$ where $\bar{k}$ is an algebraic closure of k , and similarly for $H_c^*(X, \mathbb{Q}_\ell)$, the étale cohomology groups with compact supports; these are finite dimensional $\mathbb{Q}_\ell$ -vector spaces with a continuous action of the Galois group of $\bar{k}$ over k . All Chow groups will be considered with rational coefficients.

2. LEMMAS

In this section we prove several lemmas which will be used to prove our main theorem in the next section.

Lemma 2.1. *Let W and Z be varieties over a field k and let K be a universal domain over k . Suppose the maps $CH_i(Z) \rightarrow CH_i(Z_K)$ are isomorphisms for $i = 0, 1, \dots, r$. Then any $\alpha \in CH_d(W \times Z)$ can be written as*

$$\alpha = c \sum_{i=0}^r \sum_{j=0}^{m_i} [A_{i,j} \times B_{i,j}] + \beta$$

where $A_{i,j}, B_{i,j}$ are irreducible subvarieties of W and Z respectively of respective dimensions $d-i$ and i , β is a cycle on $W \times Z$ such that $p_1(|\beta|)$ has dimension $\leq d-r-1$ and $c \in \mathbb{Q}$.

Note that we do not assume that W and Z are smooth or even irreducible.

Proof. This follows using standard methods (originally due to S. Bloch for $r = 0$) but we give a proof for the reader's convenience.

The hypothesis implies that for any extension k' of k , the maps $CH_i(Z) \rightarrow CH_i(Z_{k'})$ is an isomorphism in the given range; in particular this holds for the function fields, say $k_1, k_2, \dots, k_m$, of the irreducible components of W . Let $\iota_l : k_l \times_k Z \rightarrow W \times Z$ be the maps induced by the inclusion of the generic points and let $\alpha_l = \iota_l^*(\alpha) \in CH_{d-d_l}$, where d_l is the dimension of the irreducible component of W corresponding to k_l and $l = 1, 2, \dots, m$. By hypothesis, if $d - d_l \leq r$, then α_l is in the image of the map $CH_{d-d_l}(Z) \rightarrow CH_{d-d_l}(Z_{k_l})$ (say $\tilde{\alpha}_l \mapsto \alpha_l$), so by taking the sum of the Zariski closures of (representatives for) the cycles $k_l \times_k \tilde{\alpha}_l$ for l such that $d - d_l \leq r$ we get a cycle γ on $W \times Z$ which is a linear combinations of terms of the form $[A \times B]$ with $\dim(A) + \dim(B) = d$, $\dim(B) \leq r$ and with A one of the irreducible components of W . By construction, γ has the property that $\alpha = \alpha' + \gamma + \beta'$ where α' is supported on $W' \times Z$ with W' is a closed subvariety of W such that $W \setminus W'$ Zariski dense in W and β' has the same support properties as those desired of β . We may then replace W by W' and α by α' and complete the proof by descending Noetherian induction. $\square$

For the remainder of this section we assume that the reader is familiar with the methods of [7], in particular the notion of proper correspondences and their action on étale cohomology from [7, Section 2].

Lemma 2.2. *Let $f : X \rightarrow Y$ be a proper dominant morphism of smooth irreducible varieties over a field k with smooth geometrically irreducible generic fibre Z . Let A, B be subvarieties*

of X mapping dominantly to Y . Then the image of the proper correspondence $[A \times_Y B]$ acting on $H_c^*(X, \mathbb{Q}_\ell)$ is contained in the image of

$$H_c^*(Y, \mathbb{Q}_\ell) \otimes [A] \subset H_c^*(Y, \mathbb{Q}_\ell) \otimes H^*(X, \mathbb{Q}_\ell) \xrightarrow{f^* \otimes Id} H_c^*(X, \mathbb{Q}_\ell) \otimes H^*(X, \mathbb{Q}_\ell) \xrightarrow{\cup} H_c^*(X, \mathbb{Q}_\ell) .$$

Proof. Using the inclusion of A in X and the morphism f we may view A as giving a proper correspondence in $X \times Y$ and similarly we may view B as a proper correspondence in $Y \times X$. $[A \times_Y B]$ is then the composite of these two correspondences. It is clear from the definitions that the image of the first correspondence (mapping $H_c^*(Y, \mathbb{Q}_\ell)$ to $H_c^*(X, \mathbb{Q}_\ell)$) is exactly the image of $H_c^*(Y, \mathbb{Q}_\ell) \otimes [A]$ in $H_c^*(X, \mathbb{Q}_\ell)$. The lemma then follows from the compatibility of the composition of correspondences with the action of correspondences on cohomology. $\square$

Lemma 2.3. *Let $f : X \rightarrow Y$ be a proper dominant morphism of smooth irreducible varieties over a field k with smooth geometrically irreducible generic fibre Z . For each point y of Y , let $Z_y = f^{-1}(y)$ and let $\mathcal{Y} = \text{Spec}(\mathcal{O}_{y,Y})$. Assume that for each $i = 0, 1, \dots, r$, the specialisation maps ([8, p.399])*

$$CH_i(X \times_Y (\mathcal{Y} - \{y\}) / \mathcal{Y} - \{y\}) \rightarrow CH_i(Z_y)$$

are surjective. Let Y' be an irreducible proper subvariety of Y and let A_j, B_j , $j = 1, 2$, be irreducible subvarieties of X mapping dominantly to Y' . Suppose further that $s = \dim(B_1) - \dim(Y') \leq r$. Then the proper correspondence $[A_1 \times_Y B_1] \circ [A_2 \times_Y B_2]$ on $X \times_Y X \subset X \times X$ is equivalent (in the group of proper correspondences) to a correspondence Γ supported in $X \times_Y X$ and such that the image of $|\Gamma|$ in Y is contained in a proper closed subvariety of Y' .

Proof. As in the proof of Lemma 2.2 we may view the A_j 's as proper correspondences in $X \times Y$ and the B_j 's as proper correspondences in $Y \times X$. It then suffices to show that $[B_1] \circ [A_2]$, which is a priori a correspondence in $Y \times Y$ supported in the diagonal, is in fact supported on a proper closed subset of Y' (diagonally embedded in $Y \times Y$).

Let y be the generic point of Y' . Then B_1 gives an element in $CH_s(Z_y)$ which, using the surjectivity of the specialisation maps, must be the specialisation of an element $\alpha \in CH_s(X \times_Y (\mathcal{Y} - \{y\}) / \mathcal{Y} - \{y\})$. This implies that the element which B_1 gives in $CH_s(X \times_Y \mathcal{Y} / \mathcal{Y})$ is 0: this is clear if $\mathcal{O}_{y,Y}$ is a d.v.r. and we can reduce to this case (since $\dim(\mathcal{O}_{y,Y}) \geq 1$) using functoriality of the Gysin maps of [8] for a composite of regular embeddings. Hence B_1 is rationally equivalent, in X , to a cycle β such that $f(|\beta|) \cap Y'$ is a proper closed subset of Y' . Then β may be viewed as a proper correspondence in $Y \times X$ equivalent to $[B_1]$. This then implies the claim on the support of $[B_1] \circ [A_2]$. $\square$

Lemma 2.4. *Let $f : X \rightarrow Y$ be a proper dominant morphism of smooth irreducible varieties over a field k with smooth geometrically irreducible generic fibre Z . Let $\alpha_{i,k} \in H^{2i}(X, \mathbb{Q}_\ell)$, $i = 0, 1, \dots, r$, $k = 1, \dots, b_{2i}$, be algebraic cohomology classes such that their restrictions to the generic fibre form a basis for $\oplus_{i=0}^r H^{2i}(Z, \mathbb{Q}_\ell)$. Then the map*

$$\left(\bigoplus_{i,k} H_c^*(Y, \mathbb{Q}_\ell) \right) \xrightarrow{\oplus f^* \cup \alpha_{i,k}} H_c^*(X, \mathbb{Q}_\ell)$$

is injective.

Proof. Since the $\alpha_{i,k}$'s are algebraic classes and Z is smooth and proper, we may find algebraic cohomology classes $\beta_{i,k} \in H^*(X, \mathbb{Q}_\ell)$ such that $\alpha_{i,k} \cup \beta_{i',k'}$ restricts to $\delta_{i,i'} \cdot \delta_{k,k'} \cdot [pt] \in H^{2\dim(Z)}(Z, \mathbb{Q}_\ell)$, where δ is the Kronecker delta and $[pt]$ means the cohomology class of

a point: this follows immediately from the Hard Lefschetz theorem if A is projective and one reduces the general case to this by using Chow's lemma, alterations and the projection formula.

It follows from the construction that $f_*(\alpha_{i,k} \cup \beta_{i',k'}) = \delta_{i,i'} \cdot \delta_{k,k'} \cdot [Y] \in H^0(Y, \mathbb{Q}_\ell)$. For any cohomology class $\gamma \in H_c^*(Y, \mathbb{Q}_\ell)$, it then follows from the projection formula that $f_*(f^*(\gamma) \cup \alpha_{i,k} \cup \beta_{i',k'}) = \delta_{i,i'} \cdot \delta_{k,k'} \cdot \gamma$, which proves the injectivity. $\square$

Lemma 2.5. *Let X_1, X_2 be smooth geometrically irreducible varieties over a finite field k with q elements. Let Γ be a proper correspondence in $X_1 \times X_2$ such that $\dim(p_1(|\Gamma|)) < \dim(X) - r$. Then all the eigenvalues of the geometric Frobenius acting on $\Gamma^*(H_c^*(X_2, \mathbb{Q}_\ell)) \subset H_c^*(X_1, \mathbb{Q}_\ell)$ are algebraic integers divisible by q^{r+1} .*

Proof. This is a direct generalisation of [7, Proposition 2.4] so we do not give a detailed proof. The key point is that if $\pi : W_1 \rightarrow W_2$ is a map of smooth geometrically irreducible varieties over k such that $\dim(W_1) < \dim(W_2) - r$ then all the eigenvalues of geometric Frobenius acting on $\pi_*(H_c^*(W_1, \mathbb{Q}_\ell)) \subset H_c^*(W_2, \mathbb{Q}_\ell)$ are algebraic integers divisible by q^{r+1} . $\square$

3. PROOF OF THE MAIN THEOREM

Proof of Theorem 1.1. We apply Lemma 2.1 with $\alpha = [\Delta_Z]$, the class of the diagonal in $Z \times Z$. Let $\overline{A}_{i,j}, \overline{B}_{i,j}$ denote the Zariski closures of $A_{i,j}, B_{i,j}$ in X and let $\overline{\beta}$ be the Zariski closure of the cycle β in $X \times_Y X$. It follows that Δ_X , viewed as a proper correspondence in $X \times X$, differs from $c \sum_{i=0}^r \sum_{j=0}^{m_i} [\overline{A}_{i,j} \times_Y \overline{B}_{i,j}] + \overline{\beta}$ by a proper correspondence Γ supported in $X \times_Y X$ and such that $p_1(|\Gamma|) = p_2(|\Gamma|)$ does not contain the generic point of Y . Applying Lemma 2.1 inductively to the fibres over the generic points of $p_1(|\Gamma|)$ and the induced cycles on these fibres we see that we may write

$$[\Delta_X] = c \sum_{i=0}^r \sum_{j=0}^{m_i} [\overline{A}_{i,j} \times_Y \overline{B}_{i,j}] + d \sum_{s=0}^r \sum_{t=0}^{m_s} [C_{s,t} \times_Y D_{s,t}] + \Gamma'$$

where the $C_{s,t}, D_{s,t}$ are subvarieties of X with $f(C_{s,t}) = f(D_{s,t}) \subsetneq Y$ and $\dim(D_{s,t}/f(D_{s,t})) = s$. Furthermore, Γ' is a proper correspondence supported in $X \times_Y X$ with $\dim(p_1(|\Gamma'|)) < \dim(X) - r$.

Now repeatedly using the fact that $[\Delta_x]^2 = [\Delta_x]$ as proper correspondences and Lemma 2.3, we see that $[\Delta_X]$ can be written as a sum of products of proper correspondences, all of them supported in $X \times_Y X$, such that each product contains a factor of one of the following two types: (1) $[\overline{A}_{i,j} \times_Y \overline{B}_{i,j}]$ or (2) correspondence Γ'' such that $\dim(p_1(|\Gamma''|)) < \dim(X) - r$.

We now consider the action of such products on $H_c^*(X, \mathbb{Q}_\ell)$: By Lemma 2.2 and the fact that all our correspondences are supported in $X \times_Y X$ it follows from the projection formula that the image of any product containing a factor of type (1) must be contained in the subspace of $H_c^*(X, \mathbb{Q}_\ell)$ of the form $H_c^*(Y, \mathbb{Q}_\ell) \cdot \alpha$ where α denotes the class of some algebraic cycle on X in $H^*(X, \mathbb{Q}_\ell)$. It follows from Lemma 2.5 that the eigenvalues of the geometric Frobenius acting on the image of any product with a factor of type (2) are all divisible by q^{r+1} .

Let $\alpha_{i,k} \in H^{2i}(X, \mathbb{Q}_\ell)$, $i = 0, 1, \dots, r$, $k = 1, \dots, b_{2i}$, be algebraic cohomology classes such that their restrictions to the generic fibre Z form a basis for $\oplus_{i=0}^r H^{2i}(Z, \mathbb{Q}_\ell)$; these exist because of the assumptions on the Chow groups of Z . By Lemma 2.4 we have an injective

map

$$\left(\bigoplus_{i,k} H_c^*(Y, \mathbb{Q}_\ell) \right) \xrightarrow{\oplus f^* \cup \alpha_{i,k}} H_c^*(X, \mathbb{Q}_\ell)$$

and the image of any product as above of type (1) is contained in the image of this map. The above discussion then shows that all the eigenvalues of geometric Frobenius acting on the cokernel of this map are divisible by q^{r+1} .

We conclude the proof as in that of [7, Theorem 1.1]: without loss of generality we may assume that Y has a unique rational point and then we get the desired congruence by applying the Grothendieck-Lefschetz trace formula to X . $\square$

Corollary 3.1. *Retain the hypotheses of Theorem 1.1. Then for all $y \in Y(k)$, all the eigenvalues of the geometric Frobenius acting on $H^i(Z_y, \mathbb{Q}_\ell)$ are divisible by q^{r+1} for all $i > 2r$.*

Proof. This follows immediately from the proof of the theorem using the method of Esnault in [5, Theorem 2.1]; the idea is to use Artin's vanishing theorem for the étale cohomology of affine varieties together with the long exact sequence of cohomology with compact supports. $\square$

Remark 3.2. All our arguments are essentially motivic in nature and can be easily adapted to obtain analogous results for other cohomology theories; in particular the Hodge theoretic analogues of Theorem 1.1 and Corollary 3.1 are also true. See [7], [5] for the case $r = 0$.

4. AN APPLICATION

We now give a family of examples to which our main theorem applies. The proof of the proposition below is a variant of the proof of Otwinowska [10, Corollaire 3]; the novelty here is that we consider all hypersurfaces, not just the smooth ones.

In the following proposition all the data is assumed to be defined over a universal domain K .

Proposition 4.1. *Let B be a smooth projective variety with $\text{Pic}(B) \cong \mathbb{Z}$ generated by a very ample line bundle L and such that the cycle class maps $CH_i(B) \otimes \mathbb{Q}_\ell \rightarrow H_{2i}(B, \mathbb{Q}_\ell)$ are isomorphisms for $0 \leq i \leq 2 \dim(B)$. Let d be such that for any section $s \in H^0(B, L^{\otimes d})$, $Z(s)$, the zero scheme of s , is covered by a family of projective spaces of dimension k for some $k < \dim(B)/4$ and such that L restricts to $\mathcal{O}(1)$ on each of them. Then for any s as above, the cycle class maps $CH_i(Z(s)) \otimes \mathbb{Q}_\ell \rightarrow H_{2i}(Z(s), \mathbb{Q}_\ell)$ for $0 \leq i \leq k-1$ are isomorphisms.*

Proof. Let $A = Z(s)$ and first assume that A is smooth. We recall the proof in this case, in a slightly modified form, from [10] since we need to extend it to allow for singularities: Let G be the Grassmannian of k -planes in $\mathbb{P}(H^0(B, L))$. The hypotheses, along with the use of alterations [3], imply that we may find a smooth projective variety F with a morphism to G such that there is a dominant generically finite morphism $q : P \rightarrow A$, with $p : P \rightarrow F$ the pullback of the universal bundle of k -planes on G to F .

Let $h = c_1(\mathcal{O}_P(1)) \in CH^1(P)$. Then for any i we have

$$(1) \quad CH_i(P) = p^*CH_{i-k}(F) \oplus \left(\bigoplus_{j=1}^k h^j \cdot p^*CH_{i-k+j}(F) \right).$$

For $Z \in CH_i(A)$ we may write

$$q^*(Z) = Z_0 + h \cdot Z_1 + \cdots + h^k \cdot Z_k$$

with $Z_j \in p^*CH_{i-k+j}(F)$.

Let $H = c_1(L|_A) \in CH^1(A)$, so $h = q^*H$. Suppose $i < k$ and $Z \in CH_i(A)$ is homologically trivial. Then in the above decomposition $Z_0 = 0$ and all the Z_j 's are homologically trivial. By the projection formula, $q_*(h \cdot Z_j) = H \cdot q_*(Z_j)$ which is zero by the self-intersection formula, since, by the assumptions on B , $i_*(q_*(Z_j)) = 0 \in CH_{i+j}(B)$. Thus $q_*(q^*(Z)) = 0$ which implies that $Z = 0$ since we are working with $\mathbb{Q}$ coefficients. The isomorphisms in the theorem now follow because of the assumptions on B and the Lefschetz theorems.

Now we assume that A is reduced and irreducible but not necessarily smooth. We may find F as above, but we do not in general have a map $q^* : CH(A) \rightarrow CH(P)$ such that q^* sends homologically trivial cycles to homologically trivial cycles and q_*q^* is multiplication by the degree of q . Nevertheless, $q_* : CH(P) \rightarrow CH(A)$ is surjective so for any Z in $CH_i(A)$ with $i < k$ we may find $Z_j \in p^*CH_{i-k+j}(P)$ such that

$$Z = q_*(Z_1) \cdot H + \cdots + q_*(Z_k) \cdot H^k.$$

By the self-intersection formula as above, the image of $CH_{i+j}(A) \xrightarrow{\cdot H^j} CH_i(A)$ factors through $CH_{i+j}(A)/\equiv_{hom}$. By the assumptions on B and Artin's theorem on the cohomology of affine varieties [1, Corollaire 3.2, p.160] it follows that the maps $H_{2j}(A, \mathbb{Q}_\ell) \rightarrow H_{2j}(B, \mathbb{Q}_\ell)$ are isomorphisms for $0 \leq j < 2k$.

Now suppose that A' is a smooth hypersurface in B specialising to A , i.e. there is a commutative diagram

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{\quad} & \mathbb{P}(H^0(B, L^d)) \times_K Spec(R) \\ & \searrow & \swarrow \\ & Spec(R) & \end{array}$$

where R is a dvr with residue field K , $\mathcal{A} \rightarrow Spec(R)$ is a flat proper scheme with geometric generic fibre isomorphic to A' and special fibre isomorphic to A and the horizontal arrow is a closed embedding; such a diagram exists since L is very ample.

Using specialisation and statement for the smooth case, it follows that the maps $CH_j(A)/\equiv_{hom} \otimes \mathbb{Q}_\ell \rightarrow H_{2j}(A, \mathbb{Q}_\ell)$ are surjective, hence isomorphisms, for all $0 \leq j < 2k$, which implies that the specialisation maps $CH_j(A')/\equiv_{hom} \rightarrow CH_j(A)/\equiv_{hom}$ are also isomorphisms;

By the above discussion and the fact that $c_1(L|_{A'})$ specializes to $c_1(L|_A)$, we deduce that for $0 \leq j < k$, the specialisation maps $CH_j(A') \rightarrow CH_j(A)$ are surjective, hence must be isomorphisms since $\dim(H_{2j}(A', \mathbb{Q}_\ell)) = \dim(H_{2j}(A, \mathbb{Q}_\ell))$.

For the general case, let $A = \cup_{r=1}^m A_r$, where all the A_r 's are irreducible divisors. We prove the proposition by induction on m , the case $m = 1$ being already done. Suppose $m = 2$. We may find smooth divisors A'_1 and A'_2 specialising (as above) to A_1 and A_2 respectively such that A'_1 and A'_2 intersect transversally. By Artin's theorem as before it follows that the map $H_{2j}(A'_1 \cap A'_2, \mathbb{Q}_\ell) \rightarrow H_{2j}(A'_r, \mathbb{Q}_\ell)$ is surjective for $r = 1, 2$, $0 \leq j < k$, so that $CH_j(A'_1 \cap A'_2) \rightarrow CH_j(A'_r)$ is also surjective. By specialisation (applied to $A'_1 \cap A'_2$ and $A_1 \cap A_2$) it follows that $CH_j(A_1 \cap A_2) \rightarrow CH_j(A_r)$ is also surjective. This implies that $CH_j(A_r) \rightarrow CH_j(A_1 \cup A_2)$ is also surjective. Since the analogous map on homology is an isomorphism by Artin's theorem, it follows that the above map on Chow groups is

an isomorphism, concluding the proof in this case. The case $m > 2$ follows easily from the $m = 2$ case. $\square$

We will apply the above proposition in the case of certain homogenous spaces of Picard rank one embedded into projective space via the very ample generator. For that we shall need the following:

Lemma 4.2. *Let k be an algebraically closed field and let G_n be one of the simple algebraic groups $SL_{n+1}, Sp_{2n}, SO_{2n}, SO_{2n+1}$ over k and P_n a maximal parabolic subgroup of G_n except for the case where G_n is Sp_{2n} and P_n is the stabilizer of a maximal isotropic subspace. Let $X_n = G_n/P_n$ and L_n the (very) ample generator of $\text{Pic}(X_n)$. Then X_n is covered by a family of projective spaces of dimension $m \geq (n-1)/2$ such that the restriction of L_n to each of these subspaces is $\mathcal{O}(1)$.*

Proof. This is well known and follows from the description of the X_n 's as spaces of (isotropic) linear subspaces of the defining representation of G_n . Note that in the case of the maximal isotropic subspaces for the orthogonal groups, for the embedding induced by the Plücker embedding the restriction of $\mathcal{O}(1)$ is not the generator so this case requires a slight additional argument. $\square$

We can now prove our asymptotic generalisation of the theorem of Ax–Katz.

Theorem 4.3. *Let k be finite field with q elements, let G_n be a k -form of one of the simple algebraic groups $SL_{n+1}, Sp_{2n}, SO_{2n}$ or SO_{2n+1} and let P_n be a maximal parabolic subgroup of G_n also defined over k except for the case where G_n is Sp_{2n} and P_n is the stabilizer of a maximal isotropic subspace. Let $X_n = G_n/P_n$ and L_n the ample generator of $\text{Pic}(X_n)$. For any tuple $D := (d_1, d_2, \dots, d_t)$ of positive integers and any positive integer r , there exists an integer $N(D, r)$ such that for any $n > N(D, r)$ and any $s_i \in H^0(X_n, L_n^{\otimes d_i})$, $i = 1, 2, \dots, t$*

$$|Z(s_1, s_2, \dots, s_t)(k)| \equiv \sum_{j=0}^{r-1} \dim(b_2(X_n)) q^j \pmod{q^r}$$

where $Z(s_1, s_2, \dots, s_t)$ is the subscheme of X_n given by the vanishing of all the s_i 's.

Proof. Let $K \supset k$ be a universal domain. By applying Lemma 4.2 to X_n it follows that for fixed positive integers d, s , for any $s \in H^0(X_{n,K}, L_{n,K}^{\otimes d})$, $Z(s)$ is covered by linear spaces of dimension s if $n \gg 0$, the reason being that this is true for $X_n = \mathbb{P}^n$ (see e.g. [11]). By putting $A = X_n$ in Proposition 4.1 and using the fact that for any projective homogenous space X of a linear algebraic group the cycle class map $CH(X) \otimes \mathbb{Q}_\ell \rightarrow H(X, \mathbb{Q}_\ell)$ is always an isomorphism, we conclude that the cycle class map $CH_j(Z_s) \otimes \mathbb{Q}_\ell \rightarrow H_{2j}(Z(s), \mathbb{Q}_\ell)$ is an isomorphism for $j \leq s-1$. Moreover, $H_{2j+1}(Z(s), \mathbb{Q}_\ell)$ vanishes for $j < k$ by the Lefschetz hyperplane theorem.

By taking $s = r+1$, the previous paragraph shows that we may apply Theorem 1.1 to conclude the proof for $t = 1$. Suppose $t > 1$: by the $t = 1$ case we may find an integer $N(D, r)$ so that for all $n > N(D, r)$ the desired congruence holds for all the subschemes $\cup_{j \in T'} Z(s_j)$ where $\emptyset \neq T' \subset \{1, 2, \dots, t\}$. The theorem then follows by the Inclusion–Exclusion principle. $\square$

Example 4.4. We consider a particular homogenous space for the exceptional group $G = E_8$; our notation below is that of Bourbaki [2, p. 268]. Let B be a Borel subgroup of G and let $P \supset B$ be the parabolic subgroup of G corresponding to (removing) the root α_1 ; this

is the stabilizer (upto scalars) of the highest weight vector v in the representation of G with fundamental weight ϖ_i . Let $X = G/P$; using the data in *op. cit.* one easily sees that $\dim(X) = 78$. Let $P' \supset B$ be the parabolic subgroup of G corresponding to removing the root α_2 . By looking at the Dynkin diagram of E_8 one sees that the semi-simple part G' of a Levi subgroup of P' is isomorphic to SL_7 and one checks using the information in *loc. cit.* that the action of G' on v generates the standard representation of SL_7 . We thus see that X is covered by a family of 6-dimensional projective spaces such that the ample generator L of $\text{Pic}(X)$ restricts to $\mathcal{O}(1)$.

By proceeding as in the above theorem but using the explicit bound above we get (where we now work over a finite field k with q elements):

(1) If $s \in H^0(X, L)$ then

$$|Z(s)(k)| \equiv 1 + \sum_{i=1}^4 b_{2i}(X)q^i \pmod{q^5}$$

(2) If $s \in H^0(X, L^{\otimes 2})$ then

$$|Z(s)(k)| \equiv 1 + q \pmod{q^2}.$$

This uses the fact that a quadric in $\mathbb{P}^6$ is covered by linearly embedded $\mathbb{P}^2$'s.

Remark 4.5. One can probably use the method of Paranjape [11] or Esnault–Levine–Viehweg [6] to prove a version of Proposition 4.1 for all low degree intersections (instead of just hypersurfaces). We have not done so for the sake of brevity since the proof is not likely to require any new ideas.

REFERENCES

- [1] *Théorie des topos et cohomologie étale des schémas. Tome 3*, Springer-Verlag, Berlin, 1973. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de P. Deligne et B. Saint-Donat, Lecture Notes in Mathematics, Vol. 305.
- [2] N. BOURBAKI, *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines*, Actualités Scientifiques et Industrielles, No. 1337, Hermann, Paris, 1968.
- [3] A. J. DE JONG, *Smoothness, semi-stability and alterations*, Inst. Hautes Études Sci. Publ. Math., (1996), pp. 51–93.
- [4] H. ESNAULT, *Varieties over a finite field with trivial Chow group of 0-cycles have a rational point*, Invent. Math., 151 (2003), pp. 187–191.
- [5] ———, *Appendix to: “Congruences for rational points on varieties over finite fields” [Math. Ann. **333** (2005), no. 4, 797–809; mr 2195144] by N. Fakhruddin and C. S. Rajan*, Math. Ann., 333 (2005), pp. 811–814.
- [6] H. ESNAULT, M. LEVINE, AND E. VIEHWEG, *Chow groups of projective varieties of very small degree*, Duke Math. J., 87 (1997), pp. 29–58.
- [7] N. FAKHRUDDIN AND C. S. RAJAN, *Congruences for rational points on varieties over finite fields*, Math. Ann., 333 (2005), pp. 797–809.
- [8] W. FULTON, *Intersection theory*, Springer-Verlag, Berlin, second ed., 1998.
- [9] N. M. KATZ, *On a theorem of Ax*, Amer. J. Math., 93 (1971), pp. 485–499.
- [10] A. OTWINOWSKA, *Remarques sur les groupes de Chow des hypersurfaces de petit degré*, C. R. Acad. Sci. Paris Sér. I Math., 329 (1999), pp. 51–56.
- [11] K. H. PARANJAPE, *Cohomological and cycle-theoretic connectivity*, Ann. of Math. (2), 139 (1994), pp. 641–660.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD,
MUMBAI 400005, INDIA

E-mail address: `naf@math.tifr.res.in`